

The decay $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ and the role of σ -meson

A. Gokalp ^{*} and O. Yilmaz [†]

Physics Department, Middle East Technical University, 06531 Ankara, Turkey

(February 8, 2008)

Abstract

We study the radiative decay $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ within the framework of a phenomenological approach in which the contributions of σ -meson and ω -meson intermediate states and of the pion-loops are considered. We conclude that the σ -meson amplitude provides the dominant contribution.

PACS numbers: 12.20.Ds, 13.40.Hq

Typeset using REVTeX

^{*}agokalp@metu.edu.tr

[†]oyilmaz@metu.edu.tr

The radiative decays of neutral vector mesons into a single photon and a pair of neutral pseudoscalar mesons have been a subject of continuous interest theoretically and are becoming an area of active experimental research with the advent of high-luminosity, low energy e^+e^- -machines such as Novosibirsk and Frascati that will make possible studying these rare decays with very small branching ratios. Such measurements may serve as tests for the theoretical ideas about the interesting mechanisms of these decays because of the absence of charged particles hence the bremsstrahlung radiation, as well as shedding light on the structure of the intermediate states involved in these decays.

The decay $\rho^0 \rightarrow \pi^0\pi^0\gamma$ among other radiative ρ -meson decays was first studied by Singer [1] assuming that this decay proceeds through an $(w\pi)$ intermediate state as $\rho^0 \rightarrow (w)+\pi^0 \rightarrow (\pi^0\gamma) + \pi^0$. Renard [2] later considered σ -meson intermediate state in addition to this intermediate vector meson dominance (VDM) mechanism and noted the dependence of the decay rate as well as the photon spectrum on σ -meson mass.

The radiative decay processes of the type $V^0 \rightarrow P^0P^0\gamma$ where P and V belong to the lowest multiplets of vector (V) and pseudoscalar (P) mesons were studied by Fajfer and Oakes [3] using a low energy effective Lagrangian approach with gauged Wess-Zumino terms. They neglected scalar meson contributions and obtained the branching ratio as $B(\rho^0 \rightarrow \pi^0\pi^0\gamma)=2.89 \times 10^{-5}$. The contribution of intermediate vector mesons to the decays $V^0 \rightarrow P^0P^0\gamma$ was later considered by Bramon et al. [4] using standard Lagrangians obeying the SU(3)-symmetry. Their results were in agreement with those of Renard [2] when they were corrected using the then present-day data, however they disagree with the numerical predictions of Fajfer and Oakes [3] even if the initial expressions for the Lagrangians were the same. Their results for the decay rate and the branching ratio were $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)=1.6$ KeV and $B(\rho^0 \rightarrow \pi^0\pi^0\gamma)=1.1 \times 10^{-5}$, respectively. They also noted that final-state interactions could lead to a larger value for the decay rate $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ through the chain $\rho^0 \rightarrow \pi^+\pi^-\gamma \rightarrow \pi^0\pi^0\gamma$. Bramon et al. [5] later considered these decays within the framework of chiral effective Lagrangians and using chiral perturbation theory they calculated the decay rates for various decays of the type $V^0 \rightarrow P^0P^0\gamma$ at the one-loop level, including both

$\pi\pi$ and $K\bar{K}$ intermediate loops. They showed that the one-loop contributions are finite and to this order no counterterms are required. In this approach, the decay $\rho^0 \rightarrow \pi^0\pi^0\gamma$ proceeds mainly through charged-pion loops and Bramon et al. [5] obtained the contribution of charged-pion loops to the decay rate as $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)=1.42$ KeV which was of the same order of magnitude as the VDM contribution calculated by Bramon et al. [4] earlier. They also noted that the charged-kaon loops gave a contribution to this decay rate which was 10^3 times smaller than that of charged-pion loops. They obtained the global decay rate and the branching ratio by considering the VDM amplitude and the charged-pion loop amplitude as $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)=3.88$ KeV and $B(\rho^0 \rightarrow \pi^0\pi^0\gamma)=26 \times 10^{-6}$, respectively, and moreover noted that the photon spectrum peaked at high photon energy E_γ .

Recently, radiative decays of ρ^0 and ϕ mesons were investigated by Marco et al. [6] using the techniques of chiral unitary theory developed earlier by Oller et al. [7]. They noted that the energies of the two meson systems in the final states of these decays are too large for these decays to be treated with standard chiral perturbation theory, and using a chiral unitary approach they included the final state interaction of two mesons by summing the pion-loops through the Bethe-Salpeter equation. They obtained the branching ratio for the decay $\rho^0 \rightarrow \pi^0\pi^0\gamma$ as $B(\rho^0 \rightarrow \pi^0\pi^0\gamma)=1.4 \times 10^{-5}$, and noted that this result could be interpreted as $\rho^0 \rightarrow \gamma\sigma(\pi^0\pi^0)$ since $\pi^0\pi^0$ interaction is dominated by the σ -pole in the relevant energy regime of this decay. Furthermore, they also pointed out that if the product of the coupling constants F_V and G_V is chosen as negative $F_V G_V < 0$, where the coupling constant G_V is deduced from $\rho^0 \rightarrow \pi^+\pi^-$ decay and the coupling constant F_V from $\rho^0 \rightarrow e^+e^-$, then the result for the branching ratio becomes $B(\rho^0 \rightarrow \pi^0\pi^0\gamma)=1.0 \times 10^{-4}$; therefore the measurement of this branching ratio may serve as a test for the sign of the $F_V G_V$ product.

In this work, we follow a phenomenological approach and attempt to calculate the decay rate for the decay $\rho^0 \rightarrow \pi^0\pi^0\gamma$ by considering the σ -pole and w -pole diagrams as well as the charged-pion loops. We use the coupling constant $g_{\rho\sigma\gamma}$ that we estimated [8] using the experimental value of the branching ratio for the radiative decay $\rho^0 \rightarrow \pi^+\pi^-\gamma$. Since the

coupling constant $g_{\rho\sigma\gamma}$ was estimated in terms of the σ -meson parameters M_σ and Γ_σ , we obtain the decay rate $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ as a function of these parameters and for both signs of $g_{\rho\sigma\gamma}$.

Our calculation is based on the Feynman diagrams shown in Fig. 1. We show the amplitude for σ -meson intermediate state in Fig. 1(a). We describe the $\rho\sigma\gamma$ -vertex by the effective Lagrangian [9]

$$\mathcal{L}_{\rho\sigma\gamma}^{int.} = \frac{e}{M_\rho} g_{\rho\sigma\gamma} [\partial^\alpha \rho^\beta \partial_\alpha A_\beta - \partial^\alpha \rho^\beta \partial_\beta A_\alpha] \sigma \quad , \quad (1)$$

which also defines the coupling constant $g_{\rho\sigma\gamma}$. For the $\sigma\pi\pi$ -vertex we use the effective Lagrangian [10]

$$\mathcal{L}_{\sigma\pi\pi}^{int} = \frac{1}{2} g_{\sigma\pi\pi} M_\sigma \vec{\pi} \cdot \vec{\pi} \sigma \quad . \quad (2)$$

The decay width of the σ -meson that follows from this effective Lagrangian is given as

$$\Gamma_\sigma \equiv \Gamma(\sigma \rightarrow \pi\pi) = \frac{g_{\sigma\pi\pi}^2}{4\pi} \frac{3M_\sigma}{8} \left[1 - \left(\frac{2M_\pi}{M_\sigma} \right)^2 \right]^{1/2} . \quad (3)$$

In our calculation of the invariant amplitude, in the σ -meson propagator in Fig. 1(a) we make the replacement $M_\sigma \rightarrow M_\sigma - \frac{1}{2}i\Gamma_\sigma$, where Γ_σ is given by Eq. (3). Since the experimental candidate for σ -meson $f_0(400-1200)$ has a width (600-1000) MeV [11], in our previous work [8], using the experimental value of the branching ratio for the radiative decay $\rho^0 \rightarrow \pi^+\pi^-\gamma$ we estimated the coupling constant $g_{\rho\sigma\gamma}$ for a set of values of σ -meson parameters M_σ and Γ_σ . In our present work, we use these values of the coupling constant $g_{\rho\sigma\gamma}$. The $\omega\rho\pi$ -vertex in Fig. 1(b) is described by the effective Lagrangian [12]

$$\mathcal{L}_{\omega\rho\pi}^{int.} = \frac{g_{\omega\rho\pi}}{M_\omega} \epsilon^{\mu\nu\alpha\beta} \partial_\mu \omega_\nu \partial_\alpha \rho_\beta \pi \quad . \quad (4)$$

The coupling constant $g_{\omega\rho\pi}$ can then be determined using Vector Meson Dominance and current-field identities [12] as $g_{\omega\rho\pi}^2 \simeq 84$. The $\omega\pi\gamma$ -vertex in Fig. 1(b) is described by the effective Lagrangian [12]

$$\mathcal{L}_{\omega\pi\gamma}^{int.} = \frac{e}{M_\omega} g_{\omega\pi\gamma} \epsilon^{\mu\nu\alpha\beta} \partial_\mu \omega_\nu \partial_\alpha A_\beta \pi \quad . \quad (5)$$

The coupling constant $g_{\omega\pi\gamma}$ is then obtained from the experimental partial width [12] of the radiative decay $\omega \rightarrow \pi^0\gamma$ as $g_{\omega\pi\gamma}^2 = 3.315$. The $\rho\pi\pi$ -vertex in Fig. 1(c-e) is described by the effective Lagrangian [10]

$$\mathcal{L}_{\rho\pi\pi}^{int.} = g_{\rho\pi\pi} \vec{\rho}_\mu \cdot (\partial^\mu \vec{\pi} \times \vec{\pi}) \quad . \quad (6)$$

The experimental decay width [11] then yields the value $\frac{g_{\rho\pi\pi}^2}{4\pi} = 2.91$ for the coupling constant $g_{\rho\pi\pi}$. We describe the π^4 -vertex in Fig. 1(c-e) by the effective Lagrangian

$$\mathcal{L}^{int.} = \frac{\lambda}{4} (\vec{\pi} \cdot \vec{\pi})^2 \quad . \quad (7)$$

We note that this effective interaction results in only isospin $I=0$ amplitudes. The small $I=2$ amplitudes were also neglected in previous calculations [6] within the framework of chiral Lagrangians utilizing chiral unitary theory. We choose the coupling constant λ as $\lambda = -\frac{g_{\pi NN}^2}{4} \frac{M_\sigma^2 - M_\pi^2}{M_N^2}$ as it is obtained in the σ -model with spontaneous symmetry breaking [13] and we use $\frac{g_{\pi NN}^2}{4\pi} = 14$. We note that our effective Lagrangians $\mathcal{L}_{\sigma\pi\pi}^{int.}$ and $\mathcal{L}_{\rho\pi\pi}^{int.}$ are the ones that result from an extension of the σ model to include the isovector ρ through a Yang-Mills local gauge theory based on isospin with the vector meson mass generated through the Higgs mechanism [14]. Finally, we note that the last of the loop diagrams in Fig. 1(c-e) results from minimal coupling and ensures gauge invariance of the pion-loop amplitudes. Similar loop integrals were evaluated by Lucio and Pestiau [15] using dimensional regularization and their calculations were confirmed by Close et al. [16]. Oller [17] used their result in his calculation of the rate of the radiative decay $\phi \rightarrow K^0 \bar{K}^0 \gamma$ through an intermediate $K^+ K^-$ -loop. In our case the contribution of the pion-loop amplitudes corresponding to diagrams in Fig. 1(c-e) becomes

$$M_\pi = -\frac{e g_{\rho\pi\pi} \lambda}{2\pi^2 M_\pi^2} I(a, b) [(p \cdot k)(\epsilon \cdot u) - (p \cdot \epsilon)(k \cdot u)] \quad (8)$$

with $a = M_\rho^2/M_\pi^2$, $b = (p-k)^2/M_\pi^2$, $p(u)$ and $k(\epsilon)$ being the momentum (polarization vector) of ρ -meson and photon, respectively, and

$$I(a, b) = \frac{1}{2(a-b)} - \frac{2}{(a-b)^2} \left[f\left(\frac{1}{b}\right) - f\left(\frac{1}{a}\right) \right] + \frac{a}{(a-b)^2} \left[g\left(\frac{1}{b}\right) - g\left(\frac{1}{a}\right) \right] \quad (9)$$

where

$$\begin{aligned}
f(x) &= \begin{cases} -\left[\arcsin\left(\frac{1}{2\sqrt{x}}\right)\right]^2, & x > \frac{1}{4} \\ \frac{1}{4}\left[\ln\left(\frac{\eta_+}{\eta_-}\right) - i\pi\right]^2, & x < \frac{1}{4} \end{cases} \\
g(x) &= \begin{cases} (4x-1)^{\frac{1}{2}} \arcsin\left(\frac{1}{2\sqrt{x}}\right), & x > \frac{1}{4} \\ \frac{1}{2}(1-4x)^{\frac{1}{2}} \left[\ln\left(\frac{\eta_+}{\eta_-}\right) - i\pi\right], & x < \frac{1}{4} \end{cases} \\
\eta_{\pm} &= \frac{1}{2x} \left[1 \pm (1-4x)^{\frac{1}{2}}\right]. \tag{10}
\end{aligned}$$

In terms of the invariant amplitude $\mathcal{M}(E_\gamma, E_1)$, the differential decay probability for $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ for an unpolarized ρ^0 -meson at rest is then given as

$$\frac{d\Gamma}{dE_\gamma dE_1} = \frac{1}{(2\pi)^3} \frac{1}{8M_\rho} |\mathcal{M}|^2, \tag{11}$$

where E_γ and E_1 are the photon and pion energies respectively. We perform an average over the spin states of ρ^0 -meson and a sum over the polarization states of the photon. The decay width $\Gamma(\rho \rightarrow \pi^0 \pi^0 \gamma)$ is then obtained by integration

$$\Gamma = \frac{1}{2} \int_{E_{\gamma,min.}}^{E_{\gamma,max.}} dE_\gamma \int_{E_{1,min.}}^{E_{1,max.}} dE_1 \frac{d\Gamma}{dE_\gamma dE_1} \tag{12}$$

where now the factor $(\frac{1}{2})$ is included because of the $\pi^0 \pi^0$ in the final state. The minimum photon energy is $E_{\gamma,min.} = 0$ and the maximum photon energy is given as $E_{\gamma,max.} = (M_\rho^2 - 4M_\pi^2)/2M_\rho = 338$ MeV. The maximum and minimum values for pion energy E_1 are given by

$$\begin{aligned}
&\frac{1}{2(2E_\gamma M_\rho - M_\rho^2)} [-2E_\gamma^2 M_\rho + 3E_\gamma M_\rho^2 - M_\rho^3 \\
&\quad \pm E_\gamma \sqrt{(-2E_\gamma M_\rho + M_\rho^2)(-2E_\gamma M_\rho + M_\rho^2 - 4M_\pi^2)}] .
\end{aligned}$$

We present the results of our calculation in Table 1. Since we use the coupling constant $g_{\rho\sigma\gamma}$ that we estimated using the experimental decay rate for the radiative decay $\rho^0 \rightarrow \pi^+ \pi^- \gamma$ for a set of values of σ -meson parameters M_σ and Γ_σ in our previous work [8], we also obtain the decay rate $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma)$ as a function of these parameters. We also consider for σ meson parameters the values suggested by two recent experiments, that is $M_\sigma = 555$ MeV

$\Gamma_\sigma = 540$ MeV from CLEO [18], and $M_\sigma = 478$ MeV $\Gamma_\sigma = 324$ MeV from Fermilab E791 [19]. Since in our previous work we obtained a quadric equation for the coupling constant $g_{\rho\sigma\gamma}$ which resulted in two values of this coupling constant, one being positive and one being negative, for a given set of σ -meson parameters M_σ and Γ_σ we obtain two values for the decay rate $\rho^0 \rightarrow \pi^0\pi^0\gamma$, one using the positive value of the coupling constant $g_{\rho\sigma\gamma}$ and one using the negative value. We therefore note that the measurement of the decay rate $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ may serve as a test for the sign of this coupling constant. This result was also noticed by Marco et al. [6] for the sign of the product of the coupling constants $F_V G_V$ in their work. We further note that the measurement of the decay rate may also help to limit the ranges of the σ -meson parameters M_σ and Γ_σ as well.

We like to stress that within the framework of our phenomenological approach the main contribution to the decay rate $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ comes from the σ -meson intermediate state amplitude. Indeed, for $M_\sigma=478$ MeV, $\Gamma_\sigma=324$ MeV, and $g_{\rho\sigma\gamma}=5.92$, we obtain the value of the decay rate as $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma) = 289$ KeV and the contributions of the σ -meson intermediate state amplitude, the w -meson intermediate state amplitude and the pion-loop amplitudes are

$$\begin{aligned}\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)_\sigma &= 323 \text{ KeV}, & \Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)_\omega &= 1.46 \text{ KeV}, \\ \Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)_\pi &= 2.7 \text{ KeV}\end{aligned}\tag{13}$$

respectively. Our results for the w -meson intermediate state and pion-loop amplitudes are in reasonable agreement with the calculations of Bramon et al. [5] and Marco et al. [6]. The apparent discrepancy between our result for pion-loop contribution and that of Marco et al. [6] can be resolved by noting that the coupling constant $g_{\rho\pi\pi}$ in our work corresponds to the constant $\frac{G_V M_\rho}{f_\pi^2}$ in their work using the framework of chiral Lagrangians. They used the values $G_V = 55$ MeV, $f_\pi = 93$ MeV resulting in the coupling constant $g_{\rho\pi\pi}=4.9$. On the other hand, in our phenomenological approach we determine this coupling constant from the experimental decay rate of $\rho^0 \rightarrow \pi^+\pi^-$ [11] using the effective Lagrangian given in Eq. 6 as $g_{\rho\pi\pi}=6.0$. If we use this coupling constant as $g_{\rho\pi\pi}=4.9$ corresponding to the coupling

constants in the work of Marco et al. [6], for the contribution of the pion-loops we obtain the result $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma)_\pi = 2.14$ KeV which compares well with the result $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma)_\pi = 2.11$ KeV of Marco et al. [6]. However, since our result for the contribution of the σ -meson amplitude is much larger than the contributions of VDM and pion-loop amplitudes, our calculation results in a larger value for the decay rate $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma)$ than the previous calculations which did not include the contribution coming from the σ -meson amplitude.

Moreover, if we use the values for σ -meson parameters the values $M_\sigma=500$ MeV, $\Gamma_\sigma=250$ MeV as used by Soyeur [20], and furthermore use the value of the coupling constant $g_{\rho\sigma\gamma}=2.71$ which Titov et al. [9] obtained from Friman and Soyeur [12] analysis of ρ -meson photoproduction using VMD we obtain the decay rate as $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma) = 56.2$ KeV, or if we use $M_\sigma=500$ MeV, $\Gamma_\sigma=500$ MeV and $g_{\rho\sigma\gamma}=2.71$ we then obtain it as $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma) = 31$ KeV. In both of these calculations we determine the coupling constant $g_{\rho\pi\pi}$ from the experimental decay rate of $\rho^0 \rightarrow \pi^+ \pi^-$ which in the former case results in $g_{\rho\pi\pi}=4.5$ and in the latter case in $g_{\rho\pi\pi}=6.4$. We like to stress, however, that in our phenomenological approach, the results of which we present in Table 1, we try to estimate the coupling constants we use in our calculation utilizing the experimental information, and for the coupling constant $g_{\rho\sigma\gamma}$ we again try to relate it to σ -meson parameters through the decay rate $\Gamma(\rho^0 \rightarrow \pi^+ \pi^- \gamma)$ [8]. We also use for σ -meson parameters M_σ and Γ_σ the experimental numbers of the f_0 (400-1200) meson [11], which is the candidate for σ -meson, as summarized by Törnqvist [21].

The photon spectra for the decay rate of the decay $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$, which may be tested in future experiments, are plotted in Fig. 2 as a function of photon energy E_γ . The parameters of the σ -meson adapted for the numerical calculations are the same as above, that is $M_\sigma=478$ MeV, $\Gamma_\sigma=324$ MeV, $g_{\rho\sigma\gamma} = 5.92$. It is clearly seen that the contributions of the σ -term is much larger than the contributions of the w -term, pion-loop term as well as the interference terms. The dominant σ -term characterizes the photon spectrum which peaks at high photon energies. For comparison, we also plot the photon spectra for the same set of σ -meson parameters but for the negative value of the coupling constant $g_{\rho\sigma\gamma} = -4.49$ in Fig. 3. The general shape of the spectrum, its peaking at high photon energy as well as

the relative contributions of different terms are the same quantitatively as for the case with the positive value of the coupling constant $g_{\rho\sigma\gamma}$ shown in Fig. 2. These figures clearly show the importance of the interference terms between σ -amplitude and pion-loop amplitude and also between σ -amplitude and ω -amplitude. These interference terms change sign when we use negative value of the coupling constant $g_{\rho\sigma\gamma}$ instead of its positive value and this effect is also evident in the figures. Therefore, the measurement of the decay rate $\Gamma(\rho^0 \rightarrow \pi^0\pi^0\gamma)$ may clarify the nature of the σ -meson by limiting the range of its parameters as well as determining the sign of the coupling constant $g_{\rho\sigma\gamma}$ which plays a crucial role in analyzing photoproduction reactions within the framework of meson-exchange mechanisms such as the photoproduction of vector mesons on nucleons near threshold.

ACKNOWLEDGMENTS

We thank M. P. Rekalo for suggesting this problem to us and for his interest during the course of our work.

REFERENCES

- [1] P. Singer, Phys. Rev. **130** (1963) 2441; **161** (1967) 1694.
- [2] S. M. Renard, Nuovo Cim. **62 A** (1969) 475.
- [3] S. Fajfer and R. J. Oakes, Phys. Rev. **D 42** (1990) 2392.
- [4] A. Bramon, A. Grau and G. Pancheri, Phys. Lett. **B 283** (1992) 416.
- [5] A. Bramon, A. Grau and G. Pancheri, Phys. Lett. **B 289** (1992) 97.
- [6] E. Marco, S. Hirenzaki, E. Oset and H. Toki, Phys. Lett. **B 470** (1999) 20.
- [7] J. A. Oller, E. Oset and J. R. Pelaez, Phys. Rev. Lett. **80** (1998) 3482.
- [8] A. Gokalp and O. Yilmaz, Phys. Rev. **D 62** (2000) 093018.
- [9] A. I. Titov, T. -S. H. Lee, H. Toki and O. Streltrova, Phys. Rev.**C 60** (1999) 035205.
- [10] B. D. Serot and J. D. Walecka, in Advances in Nuclear Physics, edited by J. W. Negele and E. Vogt, **Vol. 16** (1986).
- [11] Review of Particle Physics, Eur. Phys. J.**C 3** (1998) 1.
- [12] B. Friman and M. Soyeur, Nucl. Phys. **A 600** (1996) 477.
- [13] M. A. Nowak, M. Rho and I Zahed, Chiral Nuclear Dynamics, World Scientific 1996.
- [14] B. D. Serot and J. D. Walecka, Acta Phys. Pol **B 23** (1992) 655.
- [15] J. Lucio and J. Pestiau, Phys. Rev. **D 42** (1990) 3253; **D 43** (1991) 2446.
- [16] F. E. Close, N. Isgur and S. Kumano, Nucl. Phys. **B 389** (1993) 513.
- [17] J. A. Oller, Phys. Lett. **B 426** (1998) 7.
- [18] D. M. Asner et al. (CLEO collaboration), Phys. Rev. **D 61** (1999) 012002.
- [19] E. M. Aitala et al. (E791 collaboration), hep-ex/0007028.

- [20] M. Soyeur, nucl-th/0003047.
- [21] N. A. Törnqvist, hep-ph/0008135.

TABLES

TABLE I. The decay rate $\Gamma(\rho^0 \rightarrow \pi^0 \pi^0 \gamma)$ for different σ -meson parameters

M_σ (MeV)	Γ_σ (MeV)	$g_{\sigma\pi\pi}$	$g_{\rho\sigma\gamma}$	Γ_+ (KeV)	$g_{\rho\sigma\gamma}$	Γ_- (KeV)
400	500	7.66	5.27	229	-5.23	270
450	500	6.90	6.19	256	-5.33	243
500	300	4.93	6.31	311	-4.41	202
500	500	6.36	7.21	280	-5.49	224
600	400	5.03	9.28	334	-5.63	198
555	540	6.15	8.59	294	-6.00	219
478	324	5.29	5.92	289	-4.49	215

Figure Captions:

Figure 1: Feynman Diagrams for the decay $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$

Figure 2: The photon spectra for the decay width of $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ for $g_{\rho\sigma\gamma} > 0$. The contributions of different terms are indicated.

Figure 3: The photon spectra for the decay width of $\rho^0 \rightarrow \pi^0 \pi^0 \gamma$ for $g_{\rho\sigma\gamma} < 0$. The contributions of different terms are indicated.





Figure 1

Fig. 2




Fig. 3

